

Literature:

Clinical studies listed below demonstrate usage of Microdialysis monitoring in the gastrointestinal tract following surgery.

Intraperitoneal microdialysis in the postoperative surveillance of infants undergoing surgery for congenital abdominal wall defect: A pilot study.

J Pediatr Surg. 2015 Mar 10.

Risby K, Ellebæk MB, Jakobsen MS, Husby S, Qvist N.

Mediastinal microdialysis in the diagnosis of early anastomotic leakage after resection for cancer of the esophagus and gastroesophageal junction.

Am J Surg. 2014 Jan 16.

Ellebæk M, Qvist N, Frstrup C, Mortensen MB.

Identification of anastomotic leakage after colorectal surgery using microdialysis of the peritoneal cavity.

Tech Coloproctol. 2013 Apr 30

Daams F, Wu Z, Cakir H, Karsten TM, Lange JF.

Intra-peritoneal Microdialysis and Intra-abdominal Pressure after Endovascular Repair of Ruptured Aortic Aneurysms.

Eur J Vasc Endovasc Surg. 2013 Mar 26

Hörner TM et al.

Intraperitoneal glycerol levels and lactate/pyruvate ratio: early markers of postoperative complications.

Scand J Gastroenterol. 2011 Mar 28.

Hörner TM, Norgren L, Jansson K.

Intraperitoneal microdialysis in the postoperative surveillance after surgery for necrotizing enterocolitis: a preliminary report.

Pedersen ME, Dahl M, Qvist N. J Pediatr Surg. 2011 Feb

Peritoneal microdialysis. Early diagnosis of anastomotic leakage after low anterior resection for rectosigmoid cancer.

Scand J Surg. 2009

Ellebæk Pedersen M, Qvist N, Bisgaard C, Kelly U, Bernhard A,

Møller Pedersen S.

Early detection of anastomotic leakage after pancreatoduodenectomy with microdialysis catheters: an observational Study.

Lindholm E, Bergmann GB, Haugaa H, Labori KJ, Yaqub S, Bjørn-beth BA, Line PD, Grindheim G, Kjøsén G, Pischke SE, Tønnessen TI,HPB (Oxford). 2021 Nov 5

Detection of early anastomotic leakage by intraperitoneal microdialysis after low anterior resection for rectal cancer: a prospective cohort study. Colorectal Dis. 2019 Dec:21


Ellebæk MB, Rahr HB, Boye S, Frstrup C, Qvist N.

Early detection of complications in pancreas transplants by microdialysis catheters, an observational feasibility study.

Kjøsén G et al, PLoS One. 2021 Mar 11:16.

Contact information

Principle of Microdialysis



MDialysis AB

MDialysis is the leading company devoted to the development, manufacturing and marketing of the Microdialysis technique.

The head office is located in Stockholm, Sweden, with a subsidiary in Boston MA, USA. MDialysis has distributors across the globe, responsible for local sales, service and support.

μ dialysis

MDialysis AB, Box 5049, SE-121 05 Stockholm, Sweden.

Tel: +46 8 470 10 20, Fax: +46-8-470 10 55

E-mail: info@mdialysis.se, www.mdialysis.com

Distributor

Microdialysis

To prevent Anastomosis Leakage after Gastrointestinal Surgery

Ref No: 805015SD 2024-11-27



μ dialysis

Microdialysis monitoring after Gastrointestinal Surgery


Gastrointestinal surgery is associated with a risk of post-operative complications such as anastomosis leakage and sepsis. Early diagnosis of complications and intervention is critical to improve outcome.

Intraperitoneal Microdialysis is a unique technique for early detection of complications following surgery. The technology is minimally-invasive, easy to handle, and can be used for several days.

The catheter is perfused with a sterile solution via a small pump attached to its inlet lumen. When inserted in the intraperitoneal space, small substances diffuse through the semipermeable membrane, in the tip of the catheter, into the perfusion fluid. This fluid, now known as dialysate, moves through the outlet lumen and into a collection microvial. The Microvials are exchanged with regular intervals and analyzed using the ISCUS^{flex} Microdialysis Analyzer.

Early detection of anastomosis leakage following Gastro Intestinal Surgery

The metabolite values in the collected sample provide a picture of the local metabolism and Microdialysis is a safe and reliable monitoring for surveillance of for example an anastomosis after colorectal surgery.




The Microdialysis Catheter is placed free floating in the intraperitoneal cavity during open surgery.

Microdialysis sampling


Microdialysis sampling is carried out by placing the sterile CE-certified Microdialysis Catheter in the Intraperitoneal space during Gastrointestinal surgery.

61/61 High Cut off Microdialysis Catheter



- For implantation in the intraperitoneal cavity or hepatic tissue during open surgery
- Available with 20 or 100k Dalton membrane
- Sterile, single use

106 Microdialysis Pump



The 106 Microdialysis Pump is dedicated for the perfusion of Microdialysis catheters with sterile isotonic perfusion fluid. It is handy and battery driven. The operating status is indicated by LED's and it operates at a fixed flow rate of 0.3 µl/min.

The Microdialysis monitoring system is light and portable and is not disturbing nursing actions or movement of the patient.

ISCUS^{flex} Microdialysis Analyzer

The ISCUS^{flex} Microdialysis Analyzer is specially designed for the handling of small Microdialysis sample volumes.

The system is unique for monitoring metabolic changes in tissues, intraperitoneal fluid and organs during surgery, intensive care and normal ward.

Biochemical markers:

- Glucose
- Glutamate
- Lactate
- Glycerol
- Pyruvate
- Urea
- LP-ratio



The ISCUS^{flex} Microdialysis Analyzer is easily operated by medical professionals. Data is displayed as trend curves for easy and fast interpretation.